

LA-UR-21-30669

Approved for public release; distribution is unlimited.

Title: White Source n-gamma Coincidence Measurements of gamma-Production
Cross Sections at LANSCE

Author(s): Kelly, Keegan John
Devlin, Matthew James
O'Donnell, John M.
Paris, Mark W.
Bennett, Eames Alexander

Intended for: Kickoff Meeting for NA-22-Funded Project

Issued: 2021-10-26

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



White Source n - γ Coincidence Measurements of γ -Production Cross Sections at LANSCE

Kickoff Meeting Team Presentation

Keegan J. Kelly
Matthew J. Devlin, John M. O'Donnell,
Mark Paris, and Eames Bennett

Outline

- Team Introductions
- Technical Presentation
 - Motivation
 - Other methods
 - Proposed method
 - Proof of Concept
 - Timeline and milestones
- Other items to discuss
 - LCP
 - TRL
 - Leveraging of existing technology
 - Reporting and Communications




Project Team Introductions

- **Keegan Kelly, PI**
 - Heavily involved in prompt fission neutron spectrum (PFNS) measurements at WNR FP15L at LANSCE with the Chi-Nu experiment
 - PI of LDRD-ECR and ER projects to investigate and develop neutron scattering measurement techniques
- **Matthew Devlin, Co-I**
 - PI of Chi-Nu experiment
 - GEANIE measurements of inelastic scattering and $(n,2n)$ reactions
- **John O'Donnell, Co-I**
 - Heavily involved in Chi-Nu PFNS measurement and analysis
 - Wrote and developed the Universal Analysis Code (UAC) used for analysis of data from Chi-Nu and neutron scattering data
- **Mark Paris, Co-I**
 - World-leading expert in evaluation of light-ion nuclear data evaluations
 - Evaluation results are fed directly into the potential ENDF/B libraries
- **Eames Bennett, PD**
 - Experience in Chi-Nu and neutron scattering measurement and analysis
 - One of the POCs for LDRD-funded neutron scattering analysis



Project Motivation: Active Interrogation

- n -Irradiate sample of interest
- Observe emitted γ spectrum





- γ -ray production is the common item of interest
- Inelastic n scattering cross sections are frequently the strongest contributors to γ -production cross sections
- Effectively no information on the correlated n - γ distributions
 - Not sufficient to just understand the scattering cross sections
- n transport is equally important for understanding the results

Measure correlated n - γ data to extract γ -emitting inelastic neutron scattering cross sections on ^{27}Al , ^{28}Si , and ^{16}O




Most of the γ -prod XS is in (n,n')



- γ -only measurements typically have poor E_n^{inc} resolution
- n -only measurements get cross section only (no γ info)
- Dual n - γ measurements are needed for cross section, γ production, and neutron transport
- Sensitive to all n - γ producing reactions






The LANSCE Facility: Pulsed White n Source



- Lujan Target
 - Moderated neutron target
 - Thermal – \approx MeV neutrons
 - Proton beam travels down
 - Flight paths perpendicular to p beam
- WNR Facility
 - Unmoderated tungsten target
 - Different flight path angles to p beam
 - Can obtain different flux shapes
 - $1.8 \mu\text{s}$ between p pulses







Properties of the Liquid Scintillator Array





- Span $\theta = 30\text{--}150^\circ$ at 15° increments
- Six angles in ϕ , $\approx 10\%$ of 4π
- PSD for n - γ separation
- ≈ 1 ns time resolution
- Allows for mapping of n - γ dist.



Properties of the Liquid Scintillator Array



Demonstration Measurement: Natural Carbon



- Use $t_\gamma - t_0$ to get E_n^{inc}
- Use $t_n - t_\gamma$ to get E_n^{out}
- Observe γ -coincident neutrons, with target and random coincidence backgrounds


Random Coincidence Backgrounds Eliminated

- Random coincidence rates derived from Poisson probabilities for *uncorrelated* detection rates[†]
 - true coincidence rate must be low
- Calculate the total probability for:
 - Detecting a γ at time t_γ
 - Not detecting n over coinc. time $t_n - t_\gamma$
 - Detecting n at time t_n

$$\begin{aligned}\text{Coinc. Rate} &= r_b = r_\gamma r_n \Delta t \\ \Rightarrow b &= \frac{\gamma n}{N_{t_0}} \\ \text{with } \gamma, n &= \text{counts}\end{aligned}$$



- Works remarkably well here, but what are the backgrounds?

[†]O'Donnell, NIMA 805 (2016) 87




Backgrounds from γ -Anticoincident Neutrons

- The elastic scattering $^{12}\text{C}(n,n)$ reaction is a likely source
- Do a simple Monte Carlo calculation for this background:
 - Sample incident neutrons from WNR FP15L flux shape
 - Calculate E_n^{out} from sample E_n^{inc} , convert to TOFs
 - Vary TOFs according to random γ timing, recover new $E_n^{inc'}$ and $E_n^{out'}$
 - Fill histogram with counts = $\sigma(E_n^{inc})$
- Possible to extract cross sections from this background?...maybe...




MCNP Simulations of Carbon Data

Data




MCNP




- Chi-Nu experiment analysis relied heavily on MCNP simulations
- MCNP can guide n - γ detector efficiencies
- Number of internal scatters can be investigated
- Understanding neutron interactions with the environment is important




Extract n , γ , and Correlated n - γ Distributions




(a)




(b)




(c)



(d)





(e)





(f)

Corr. n - γ Distributions for Wide Range of E_n^{inc}



Goal and Expected AI Yield





- Use γ -coincident n yields to extract γ -production cross sections
- Initially assume level decay branching ratios from each excited state
- Can be used to cross check data γ -anticoincident data, and n -only MCNP simulations

And now for a couple of interesting analysis details that we hope to exploit for the analysis of these data



Obtain 2D n Efficiency from $^{12}\text{C}(n,n'\gamma)$


- Typical treatments of detection efficiency work poorly for neutrons
- Need complete description of n interactions with exp. environment
 - Especially for smooth distributions (e.g., high level density scattering)
- This was handled with MCNP for Chi-Nu PFNS measurements
- Could be more accurately done with measurements



Iterative Unfolding to Isolate Excited States


From last week:
$$m_{\alpha|\beta}^{(n+1)}(E) = \frac{m_{\alpha|\beta}^{(n)}(E)c_{\alpha}(E)}{\sum_{i=1}^N \mathcal{R}(E, E_i)m_{\alpha|\beta}^{(n)}(E_i)}$$

- Clearer view of states in data
- Informs treatment of n response
- Clarifies expected γ rays
- Confirms inelastic state populations




Sensitive to *All* n - γ Producing Reactions


- Fe levels are reasonably dense
- The liquid scint. time resolution allows for ^{56}Fe low-lying state separation
- Natural Pt shows inelastic scattering, $(n,2n)$, and $(n,3n)$ reactions, with separation
- Elastic scattering data also exist from these measurements
- Potential for correlated measurements of these different cross sections



Experimental Setup for Aluminum Measurement



Project Timeline



- Proposed milestones are repetitive for each proposed target:
 - Obtain data from each target near start of each FY
 - Analyze data during same FY as measurement
 - Work towards publication near end of FY and start of next FY
 - Include data in evaluation as they are available (typically before
- Potential for final meas. to use only CLYC detectors may arise



Other Items to Discuss

- LCP, deliverables, work scope, and milestones
 - Discussion of acceptance criteria for milestones?
- Leveraging existing and developing detection and sample setups, as well as analysis techniques developed under LDRD funding
- Team consists
 - 3 more senior staff members (Co-I's)
 - 1 early career scientist (PI)
 - 1 postdoc
- Technical Readiness Level (TRL)
 - Starting: 8/10, system demo, nearing complete analysis path
 - Ending: 10/10, system complete and in production use
- Reporting requirements
- Constraints (beamtime availability)
- Risks and opportunities
- Communications plan and possible stakeholders

